Dates of revision: 2024 February

You should read this written explanation before use. Please keep it so that you can read it when you need to.

Third-Class OTC drugs

Protect the wounds with a transparent film.

Coloskin_®

Liquid Plaster

Apply Coloskin® for wounds, such as minor cuts, abrasions, hangnails, and chapped skin.

When dry, it would form a transparent film that protects the wounds from irritation of water or hot water and germs.

And Coloskin_® of which the film would closely adhere to skin, and not be easily peeled off by work to use water or wash hands, is the Liquid Plaster which does not require an ordinary bandage.

Precautions

When not to use the product (If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- 1. The medicine should not be used on the following parts.
 - 1 Erosion, purulence, bleeding wounds, and major wounds
 - ② Eyes and around eyes, face, head and mucosa (such as oral cavity and nasal cavity, etc.)
 - 3 Wounded skin by shaving or depilation.
- 2. Do not intentionally inhale as the medicine contains organic solvents.

Consultation

 The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.

Patients who have experienced allergic symptoms associated with drugs, etc.

If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately
discontinue the use, and show this document to your physician, pharmacist, or registered salesperson for a
consultation.

Areas Concerned	Symptoms
Skin	Rash/Redness, Itching, Contact dermatitis, Hot sensation, etc.

Indication

Minor cuts, Abrasions, Hangnails, and Chapped skin

Dosage and administration

Clean the affected lesion. Apply an appropriate amount of Coloskin® to the wound and dry thoroughly.

Precautions of dosage and administration


① Applying on the wounds may sting for a moment, but it will subside as soon as dry, and the film would protect

the wounds.

- 2 Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water, and immediately seek an ophthalmologist for medical treatment.
- 3 Applied area should be kept away from fire or flame.
- 4 When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.
- 5 This medicine is only for external use and should not be internal use.
- ⑥ If blood, water, or oil from cream or ointments remains on the affected area, Coloskin_® may not adhere well, so wipe the area off thoroughly before use.
- 7 Do not apply the medicine extensively around the wounds.
- 8 The film is naturally peeled off from the surrounding area in your daily life. If the wounds have not yet healed when the film peels off, apply Coloskin® again. There is no set time or frequency of use per day.

Usage

- ① Open a hole in tube with prong of cap.
- 2 Push out the tube from the fold (tail) end towards the tip.



- 3 When you are in haste to remove the film,
 - Scrape it off from the surrounding area after steaming and warming with steamed towel.
 - Wipe it off with enamel remover as dissolving.
 - Apply Coloskin® again over the film and wipe it off before dries.

Ingredient and amount per 100g

Pyroxylin · · · · · 15.95g d-Camphor · · · · · 2.8g

Additives · · · · · Benzyl alcohol, Castor oil, Ethyl acetate, Butyl acetate

Precautions for storage and handling

- ① Seal hermetically and store in a cool place away from direct sunlight.
- 2 Keep out of reach of children.
- 3 Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
- 4 Keep away from fire or flame.
- (5) Soaking the cap in hot water may be easier to open if the cap of the tube does not open. (Do not warm by direct flame.)
- ⑥ After use, wipe Coloskin® from the opening of the tube and cap well before closing the cap.
- Attention should be paid when using Coloskin_® if it gets on clothes, furniture, or floors, etc., it may be very difficult to remove and cause changes in color tone and material quality.
- 8 Do not use any medicine after the expiration date which is indicated on the box.

<Disclaimer on multilingual product information>

- •This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- •Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- •Multilingual product information is a tentative translation by our company, and may be modified or altered without notice.
- •Our company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

[Contact]

TOKYOKOSHISHA Customer Support Center

Tel: 03-3862-4081

Office Hours: 10:00 - 16:00 (Except Saturdays, Sundays, and Public Holidays)

[Manufacture and Sales]

TOKYOKOSHISHA CO., LTD.

3-10-9 Iwamoto-cho, Chiyoda-ku, Tokyo

Coloskin_® is the Liquid Plaster which protects the wounds easily and cleanly through all seasons as a household medicine.





